

Review of 2-Dimension Coordinate Geometry

'AKA... Numbers, Relations and Functions 10'

Oct 16-6:43 PM

Cartesian Plane

Associates each point with a pair of numbers (**ordered pair**).

Oct 16-6:45 PM

Calculating Slope

$a =$

#1. Graph

Slope = $\frac{\text{Rise}}{\text{Run}}$

$m = \frac{-6}{8} = -\frac{3}{4}$

#2. Two Points

ex: $(-3, 5)$ & $(1, -7)$

$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-7 - 5}{1 - (-3)} = \frac{-12}{4} = -3$

#3. Equation

ex: Determine the slope of...

$3x - 2y - 6 = 0$

$-2y = -3x + 6$

$-2 \quad -2 \quad -2$

$y = \frac{3}{2}x - 3$

Example...

Find the value for x if the line segment joining the points $(0, 0)$ and $(-2, 4)$ has a slope value of $\frac{2}{3}$

$m = \frac{y_2 - y_1}{x_2 - x_1}$

$\frac{2}{3} = \frac{4 - 0}{-2 - x}$

$\frac{2}{3} \cdot (-2 - x) = 4$

$-2(-2 - x) = 4(3)$

$+4 + 2x = 12$

$2x = 8$

$x = 4$

Oct 10-8:20 PM

Intercepts

x intercept

Where does it cross the x - axis? (Let $y = 0$)

y intercept

Where does it cross the y - axis? (Let $x = 0$)

Ex. $2x - 3y = 12$

x-int let $y=0$

$2x - 3(0) = 12$

$2x = 12$

$x = 6$

y-int let $x=0$

$2(0) - 3y = 12$

$-3y = 12$

$y = -4$

$m = \frac{4}{6} = \frac{2}{3}$

Oct 16-7:36 PM

Graphing Linear Functions

NOTES - Graphing Linear Relationships.docx

Method #1 - Table of Values (must have at least 3 points)

ex: $3x - 6y + 18 = 0$

$3(-1) - 6y + 18 = 0$
 $-3 - 6y + 18 = 0$
 $-6y + 15 = 0$
 $-6y = -15$
 $y = \frac{5}{2}$

$3(0) - 6y + 18 = 0$
 $-6y + 18 = 0$
 $-6y = -18$
 $y = 3$

x	y
-1	2.5
0	3
1	

Method #2 - Using the slope/intercept form of the equation

- put equation in the form.

$y = mx + b$ ← y-int

- plot the y intercept
- use slope = $\frac{\text{Rise}}{\text{Run}}$ to plot other points.

ex: $3x - 2y = -4$

$\frac{-2y}{-2} = \frac{-3x - 4}{-2}$
 $y = \frac{3}{2}x + 2$

Oct 10-10:06 PM


Method #3 - Using x / y intercepts

ex: $x - 5y - 10 = 0$

x-int let y=0
 $x - 5(0) - 10 = 0$
 $x - 10 = 0$
 $x = 10$

y-int let x=0
 $0 - 5y - 10 = 0$
 $-5y = 10$
 $y = -2$

Oct 10-10:18 PM



Sep 3-10:31 AM

What about vertical versus horizontal lines???

Graphs of Special Lines

- horizontal lines - slope value of zero

ex: (3, -4) & (-1, -4)

- vertical lines - slope value is **undefined**

ex: $x = 2$

Oct 4-11:44 AM

WHY WE CAN'T DIVIDE BY ZERO...

Sep 5-9:46 AM

The image shows a video player interface. The main content is a black screen with the equation $y = mx + b$ in white. Below the equation, the text 'Graph!' and 'Westerville South High School' is visible. At the bottom of the video frame, there is a progress bar showing '0:02 / 4:16'. Below the video frame, the title 'Graph! (WSHS Math Rap Song)' is displayed.

Sep 3-1:47 PM

HOMEWORK...

Puzzle Worksheet - Graphing Lines.docx

Oct 4-11:23 AM

QUESTIONS FROM THE HOMEWORK???

Sep 6-9:53 AM

WARM-UP: Let's Review... PRIOR KNOWLEDGE???

WORDS You Need to Communicate Effectively Warm Up - Prior Knowledge for Coordinate Geometry.docx


1. Match each term with the best example or description on the right.

a) linear equation	ii) the value 3 in the equation $y = 3x + 1$
b) x- and y-intercepts	iii) $\{(1, 5), (2, 6), (3, 7)\}$ in the solution set $\{(1, 5), (2, 6), (3, 7)\}$
c) slope	iv) in a relationship, the variable graphed on the y-axis
d) linear inequality	v) $2y = 3x + 7$
e) dependent variable	vi) $3 = x + 5$
f) domain	vii) term used to describe a solution set from the set of real numbers
g) range	viii) $\{(5, 6), (0, -5)\}$ for the graph of $y = 4x - 5$
h) discrete	ix) $\{(5, 6, 7)\}$ in the solution set $\{(1, 5), (2, 6), (3, 7)\}$
i) continuous	x) in a relationship, the variable graphed on the x-axis
j) independent variable	xi) term used to describe a solution set from the set of integers
k) quadrant I	xii) the part of the coordinate plane where $x > 0$ and $y > 0$

(Handwritten notes: A coordinate plane with a red line $y = x + 5$ and a purple line $y = 4x - 5$. The region where they intersect is labeled (x, y) . Marginal notes include 'viii', 'vii', 'v', 'vi', 'vii', 'viii', 'ix', 'x', 'xi', 'xii'.)

Answers

1.



⑧ $2x - 2y + 5 = 0$

$$\begin{aligned} -2y &= -2x - 5 \\ \frac{-2y}{-2} &= \frac{-2x}{-2} - \frac{5}{-2} \\ y &= 1x + 2.5 \end{aligned}$$

REVIEW OF TERMS AND CONNECTIONS

CONNECTIONS You Need for Success

Applying Number Concepts

Number Classification

When working with linear equations and linear inequalities, the domain and range may be restricted to a specific set of numbers. Knowing how the sets of numbers are different is important when interpreting and solving problems and when graphing. For example:

- If $\{(x, y) \mid x \in W, y \in W\}$, the variables x and y are from the set of whole numbers and the problem has a whole-number solution, such as $(4, 6)$. If the problem is represented graphically, the graph will be in the first quadrant.
- If $\{(x, y) \mid x \in R, y \in R\}$, the variables x and y are from the set of real numbers and the problem has real-number solutions, such as $(-4, 5.7)$ or $(\sqrt{2}, \sqrt{8})$. If the problem is represented graphically, the graph could be in any of the four quadrants.

2. Give an example of an ordered pair that could be in each solution set.

a) $\{(x, y) \mid x \in I, y \in I\}$	c) $\{(k, j) \mid k \in N, j \in N\}$
b) $\{(m, p) \mid m \in R, p \in R\}$	d) $\{(x, y) \mid x \geq 0, x \in R, y \geq 0, y \in R\}$

Answers

2.

Linear Inequalities:
 equality sign - could be one of the following...

$>$ LESS THAN
 GREATER THAN
 $<$ LESS THAN
 GREATER THAN OR EQUAL TO
 \geq LESS THAN OR EQUAL TO
 GREATER THAN OR EQUAL TO
 \leq LESS THAN OR EQUAL TO
 GREATER THAN OR EQUAL TO
 \neq NOT EQUAL TO

When solving an in-equation, all the steps are the same EXCEPT when it comes to isolating..

4 $<$ 11, fill in the box. $4 < 11$

Now divide both by -1

-4 $>$ -11, fill in the box.

RULE: If you multiply or divide by a negative, reverse the inequality sign!!!

Solving Equations	Solving Inequality
$2x - 5 = 7$ $2x = 12$ $x = 6$	$2x - 5 > 7$ $2x > 12$ $x > 6$
$4 - 3x = 19$ $-3x = 15$ $x = -5$	$4 - 3x \leq 19$ $-3x \leq 15$ $x \geq -5$

Sep 26-1:51 PM

Rearranging Linear Inequalities

ex $4x - 3y + 12 = 0$ $4x - 3y + 12 > 0$

$\frac{-3y}{-3} = \frac{-4x - 12}{-3}$

$y < \frac{4}{3}x + 4$

** new*

Jan 30-2:54 PM

Try

a) $2x + y \geq 7$
 $y \geq -2x + 7$

b) $x - 5y + 10 \leq 0$
 $-5y \leq -x - 10$
 $\frac{-5y}{-5} \geq \frac{-x - 10}{-5}$
 $y \geq \frac{1}{5}x + 2$

$\frac{x+10 \leq 15y}{\frac{1}{5}x + 2 \leq y}$

Jan 30-2:58 PM

$x > 2$ $2 < x$

*

Jan 30-3:03 PM

$$\begin{array}{l}
 4^{+2} < 11^{+2} \\
 6^{-5} < 13^{-5} \\
 1^{x10} < 8^{x10} \\
 10^{\div 2} < 80^{\div 2} \\
 5^{x(3)} < 40^{x(3)} \\
 -15 > -120
 \end{array}$$

Jan 30-2:44 PM

Try these:

① $-5 + 2x \leq 9$
 $2x \leq 14$
 $x \leq 7$

② $10 - 4x \leq -14$
 $-4x \leq -24$
 $x \geq 6$

Rearrange "y mx+b"

ex $-3x + 2y - 12 > 0$ $4x - 3y \geq 9$
 $\frac{2y}{2} > \frac{3x}{2} + \frac{12}{2}$ $-\frac{3y}{-3} \geq \frac{-4x}{-3} + \frac{9}{-3}$
 $y > \frac{3}{2}x + 6$ $y \leq \frac{4}{3}x - 3$

Jan 30-1:14 PM

5.1 Graphing Linear Inequalities in Two Variables

GOAL
Solve problems by modelling linear inequalities in two variables.

EXPLORE...

• For which inequalities is (3, 1) a possible solution? How do you know?

a) $13 - 3x > 4y$
 $13 - 3(3) > 4(1)$
 $4 > 4$
False!

b) $2y - 5 \leq x$
 $2(1) - 5 \leq 3$
 $-3 \leq 3$
True

c) $1 + 3 < 10$
 $4 < 10$
True

d) $1 \geq 9$
False

VERIFY

6.1 Page 1

NOTES - Graphing a Linear Inequation.docx

When the solution set to a linear inequality is continuous and the sign does not include equality, use a dashed line for the boundary and shade the solution region.

Example: Graph the solution to: $2x - 3y < 6$.

First, solve for the equation in the slope - y intercept form ($y = mx + b$).

$$\begin{array}{l}
 2x - 3y < 6 \\
 -3y < -2x + 6 \\
 y > (2/3)x - 2
 \end{array}$$

STEP 1: Graph the boundary line

Find the "equals" part, which is the line $y = (2/3)x - 2$. It looks like this:

But this example is a **strict inequality**. That is, it's only "y greater than." We denote strict inequalities on the number line (such as $x > 5$) by using an open dot instead of a closed dot. In the case of these linear inequalities, the notation for a strict inequality is a dashed line. So the boundary line of the solution region actually looks like this:

STEP 2: Decide on dashed or solid

By using a dashed line, we can still identify the boundary line, but the dashed line indicates that the boundary line isn't included in the solution. Since this is a "y greater than" inequality, we will shade above the line, so the solution looks like this:

STEP 3: Pick a 'test point' and verify
STEP 4: Shade or stipple

VIDEO - Graphing Inequalities

[Click HERE to watch the video!!!](#)

Oct 9-12:37 PM

WORK that NEEDS to be DONE at HOME...
 (also known as 'Homework')

- 1) READ Given Notes on Graphing Inequalities
- 2) Watch Video Link - lesson is on the website
- 3) Complete logic problem and pass it in!

DUE MONDAY...

Logic - Three Little Pigs.doc

Sep 6-9:31 AM

APPLY the Math EXAMPLE FROM TEXT P. 213

EXAMPLE 1 Solving a linear inequality graphically when it has a continuous solution set in two variables

Graph the solution set for this linear inequality:
 $-2x + 5y \geq 10$

Robert's Solution: Using graph paper

Linear equation that represents the boundary:
 $-2x + 5y = 10$

The variables represent numbers from the set of real numbers.
 $x \in \mathbb{R}$ and $y \in \mathbb{R}$

y-intercept:
 $-2x + 5y = 10$
 $-2(0) + 5y = 10$
 $5y = 10$
 $y = 2$
 The y-intercept is at (0, 2).

x-intercept:
 $-2x + 5(0) = 10$
 $-2x = 10$
 $x = -5$
 The x-intercept is at (-5, 0).

Since the linear inequality has the possibility of equality (\geq), and the variables represent real numbers, I know that the solution region includes all the points on its boundary. That's why I drew a solid green line through the intercepts.

To find out, I substituted the coordinates of a point in the half plane below the line. I used (0, 0) because it made the calculations simple. I already knew that the solution region includes points on the boundary, so I didn't need to check a point on the line.

Since my test point below the boundary was not a solution, I shaded the half plane that did not include my test point. This was the region above the boundary. I used green shading to show that the solution set belongs to the set of real numbers. Since the domain and range are in the set of real numbers, I knew that the solution set is continuous. Therefore, the solution region includes all points in the shaded area and on the solid boundary.

continuous
 A connected set of numbers in a continuous set, there is always another number between any two given numbers. Continuous variables represent things that can be measured, such as time.

solution region
 The part of the graph of a linear inequality that represents the solution set. The solution region includes points on its boundary if the inequality has the possibility of equality.

half plane
 The region on one side of the graph of a linear relation on a Cartesian plane.

Communication Tip
 If the solution set of a linear inequality is continuous and the sign includes equality (\geq or \leq), a solid green line is used for the boundary, and the solution region is shaded green.

6.1 Page 4

EXAMPLE #2:

$y < 5x + 1$

Test (3,1)
 $y < 5x + 1$
 $1 < 5(3) + 1$
 $1 < 16$

Oct 11-11:00 AM

EXAMPLE #3:


$3y \leq -5x$

$y \leq -\frac{5}{3}x$


Test Point
 $(4, 1)$
 $3(1) \leq 5(4)$
 $3 \leq 20$

$3(-3) \leq -5(-1)$
 $-9 \leq 5$
 True!

Oct 11-11:02 AM



Oct 11-11:02 AM



Sep 9-9:37 AM

HOMEWORK...

[Puzzle Worksheet - Graphing Linear Inequalities with Two Variables.pdf](#)

Oct 9-4:16 PM

WARM-UP: Graph each of the following...

1) $y \leq -\frac{8}{3}x - 3$

2) $2x + 5y - 20 > 0$

Oct 11-11:04 AM

Graphs of Linear In-Equalities

Sometimes the domain and range are stated as being in the set of integers. This means that the solution set is **discrete** and consists of separate or distinct parts. Discrete variables represent things that can be counted, such as people in a room. This means that the solution region is not shaded but rather stippled with points.


So when interpreting the solution region for a linear inequality, consider the restriction on the domain and range of the variables.

If the solution set is **continuous**, all the points in the solution region are in the solution set. (Shaded)

If the solution set is **discrete**, only specific points in the solution region are in the solution set. This is represented graphically by stippling.

Some solution sets may be restricted to specific quadrants. For example, most linear inequalities representing real-world problem situations have graphs that are restricted to the first quadrant.

Here are some examples:



Handwritten notes explaining set notation:


$$\{(x,y) | y > -2x + 5, x \in \mathbb{R}, y \in \mathbb{R}\}$$

"set of" points to the curly braces, "such that" points to the vertical bar, and "belongs to" points to the elements $x \in \mathbb{R}, y \in \mathbb{R}$.

Oct 9-3:07 PM

Jan 31-2:56 PM

Let's do a couple more...



Oct 11-11:16 AM

Try

a) $\{(x,y) | y \geq \frac{1}{4}x - 3, x \in \mathbb{I}, y \in \mathbb{I}\}$

stip

stipple here

Test (0,0)
 $0 \geq \frac{1}{4}(0) - 3$
 $0 \geq -3$
 True

Feb 3-12:48 PM

b) $\{(x, y) \mid 2x - 3y < -6, x \in \mathbb{W}, y \in \mathbb{W}\}$

$-3y < -2x - 6$
 $y > \frac{2}{3}x + 2$

Test (0,0)
 $2(0) - 3(0) < -6$
 $0 < -6$
 False

EXAMPLE 1 Solving a linear inequality graphically when it has a continuous solution set in two variables

Graph the solution set for this linear inequality:
 $-2x + 5y \geq 10$

Your Turn
 Compare the graphs of the following relations. What do you notice?
 $-2x + 5y \geq 10$ $-2x + 5y = 10$ $-2x + 5y < 10$

Answer

Feb 3-12:56 PM

6.1 Page 6

EXAMPLE 2 Graphing linear inequalities with vertical or horizontal boundaries

Graph the solution set for each linear inequality on a Cartesian plane.

a) $\{(x, y) \mid x - 2 > 0, x \in \mathbb{R}, y \in \mathbb{R}\}$
 b) $\{(x, y) \mid -3y + 6 \geq -6 + y, x \in \mathbb{I}, y \in \mathbb{I}\}$

Wynn's Solution

a) $x - 2 > 0$
 $x > 2$

The variables represent numbers in the set of real numbers.
 $x \in \mathbb{R}$ and $y \in \mathbb{R}$

The domain and range are stated as the set of real numbers. The solution set is continuous, so the solution region and its boundary will be green in my graph.

I drew the boundary of the linear inequality as a dashed green line because I knew that the linear inequality ($>$) does not include the possibility of x being equal to 2.

I needed to decide which half plane to shade. For x to be greater than 2, I knew that any point to the right of the boundary would work.

The solution region includes all the points in the shaded area because the solution set is continuous. The solution region does not include points on the boundary.

b) $-3y + 6 \geq -6 + y$
 $-4y \geq -12$
 $-y \geq -3$
 $y \leq 3$

Since the linear inequality has only one variable, y , I isolated the y .

As I rearranged the linear inequality, I divided both sides by -4 . That's why I reversed the sign from \geq to \leq .

The domain and range are stated as being in the set of integers. I knew this means that the solution set is **discrete**.

I knew that points with integer coordinates below the line $y = 3$ were solutions, so I shaded the half plane below it orange.

I knew the linear inequality includes 3, so points on the boundary with integer coordinates are also solutions to the linear inequality.

I stippled the boundary and the orange half plane with green points to show that the solution set is discrete.

The solution region includes only the points with integer coordinates in the shaded region and along the boundary.

Communication Tip
 If the solution set to a linear inequality is continuous and the sign does not include equality ($<$ or $>$), a dashed green line is used for the boundary and the solution region is shaded green.

discrete
 Consisting of separate or distinct parts. Discrete variables represent things that can be counted, such as people in a room.

Communication Tip
 If the solution set to a linear inequality is discrete, the solution region is shaded orange and stippled with green points. If the sign includes equality (\leq or \geq), the boundary is also stippled. An example of this is shown to the left. If equality is not included ($<$ or $>$), the boundary is a dashed orange line. An example of this is shown below.


6.1 Page 7

Oct 9-3:13 PM

HOMEWORK...

p. 221: #3, #4, #6

$\frac{\text{not told}}{\text{shade}}$
 $\frac{N, W, I}{\text{stipple}}$
 $\frac{R}{\text{shade}}$



Feb 4-12:41 PM

INVESTIGATE the Math

Amir owns a health-food store. He is making a mixture of nuts and raisins to sell in bulk. His supplier charges \$25/kg for nuts and \$8/kg for raisins.

1 What quantities of nuts and raisins can Amir mix together if he wants to spend less than \$200 to make the mixture?

A. Suppose that Amir wants to spend exactly \$200 to make the mixture. Work with a partner to create an equation that represents this situation.

B. To what set of numbers does the domain and range of the two variables in your equation belong? Use this information to help you graph the equation on a coordinate plane.

C. Explain why the graph is a line segment, not a ray or a line.

D. What region of the coordinate plane includes points representing quantities of nuts and raisins that Amir could use if he wants to spend less than \$200? How do you know?

E. There are many possible solutions to Amir's problem. Plot at least three points that represent reasonable solutions to Amir's problem. Explain why you chose these points.

Answers

A. Let x represent the number of kilograms of nuts, and let y represent the number of kilograms of raisins.
 $25x + 8y = 200$

B. The domain and range belong to the set of real numbers greater than or equal to 0.

Raisins and Nuts

C. The graph is a line segment because it has endpoints at the x -axis and y -axis.

D. The region below the line segment includes points representing quantities of nuts and raisins that Amir could use. Answers will vary, e.g., I picked a point in the region above and below the line and tested both algebraically:


Raisins and Nuts

E. Answers will vary, e.g., I chose these points because they are in the region below the line. I know that points in the region below the line will solve the problem. However, even though points such as (0, 0), (7, 0) and (0, 24) are below the line, they wouldn't make very good mixtures because they either involve only one quantity, or no quantities at all.

Raisins and Nuts

Test a point above the line. (8, 8), or 8 kg of nuts and 8 kg of raisins: $25x + 8y = 25(8) + 8(8) = 264$ So, 8 kg of nuts and 8 kg of raisins would cost \$264, which is more than \$200.	Test a point below the line. (5, 5), or 5 kg of nuts and 5 kg of raisins: $25x + 8y = 25(5) + 8(5) = 165$ So, 5 kg of nuts and 5 kg of raisins would cost \$165, which is less than \$200.
---	---

6.1 Page 2



Sep 11-3:02 PM

Reflecting

F. Discuss and then decide whether the **solution set** for Amir's problem is represented by

- points in the region above the line segment.
- points in the region below the line segment.
- points on the line segment.

G. Why might the line segment be considered a boundary of the solution set?

H. Why might you use a dashed line segment for this graph instead of a solid line segment?

Answers

F. **i)** Points in the region above the line segment are not part of the solution set because they result in costs greater than \$200.
ii) Points in the region below the line segment represent the solution set because they result in costs less than \$200.
iii) Points on the line segment are not part of the solution set because they result in costs that are exactly \$200.

G. The line segment might be considered a boundary because the solution set is completely on one side of it. It separates the points that are in the solution set from the points that are not.

H. A dashed line segment shows that the points on the line segment are not part of the solution set.

solution set
 The set of all possible solutions

6.1 Page 3

Applications...Apply your skills to a context

EXAMPLE #2: HANDOUT - Application of a Linear Inequality.docx

Malia and Lainey are competing in a spelling quiz. Malia gets a point for every word she spells correctly. Lainey is younger than Malia, so she gets 3 points for every word she spells correctly plus one bonus point. What combination of correctly spelled words for Malia and Lainey result in Malia spelling more? Choose two combinations that make sense and explain why.

Step 1: Declare variables $m \rightarrow \# \text{ of words for Malia}$
 $l \rightarrow \# \text{ of words for Lainey}$


Step 2: State restrictions $m \geq 0, m \in \mathbb{W}$
 $l \geq 0, l \in \mathbb{W}$

Step 3: Develop the inequation

$$m > 3l + 1$$

$$(y > 3x + 1)$$

Step 4: Graph the solution set (MUST include labels/scales)



Sep 11-2:34 PM

EXAMPLE 3 p. 218 Solving a real-world problem by graphing a linear inequality with discrete whole-number solutions

A sports store has a net revenue of \$100 on every pair of downhill skis sold and \$120 on every snowboard sold. The manager's goal is to have a net revenue of more than \$600 a day from the sales of these two items. What combinations of ski and snowboard sales will meet or exceed this daily sales goal? Choose two combinations that make sense, and explain your choices.




Step 1: Declare variables $x \rightarrow \# \text{ of skis sold}$
 $y \rightarrow \# \text{ of snowboards sold}$

Step 2: State restrictions $x \geq 0, x \in \mathbb{W}$
 $y \geq 0, y \in \mathbb{W}$

Step 3: Develop the inequation $100x + 120y \geq 600$

Step 4: Graph the solution set (MUST include labels/scales)



Sep 11-3:44 PM

EXAMPLE 3 p. 218 Solving a real-world problem by graphing a linear inequality with discrete whole-number solutions

A sports store has a net revenue of \$100 on every pair of downhill skis sold and \$120 on every snowboard sold. The manager's goal is to have a net revenue of more than \$600 a day from the sales of these two items. What combinations of ski and snowboard sales will meet or exceed this daily sales goal? Choose two combinations that make sense, and explain your choices.

let x = # of skis sold
y = # of snowboards sold

Jerry's Solution

The relationship between the number of pairs of skis, x , the number of snowboards, y , and the daily sales can be represented by the following linear inequality:
 $100x + 120y > 600$

The variables represent whole numbers.
 $x \in \mathbb{W}$ and $y \in \mathbb{W}$

$100x + 120y > 600$
 $120y > 600 - 100x$
 $120y > 600 - 100x$
 $120 > 120$
 $y > \frac{600 - 100x}{120}$
 $y > 5 - \frac{5x}{6}$
 $y > -\frac{5x}{6} + 5$

I defined the variables in this situation and wrote a linear inequality to represent the problem.

I knew that only whole numbers are possible for x and y , since stores don't sell parts of skis or snowboards. Because the domain and range are restricted to the set of whole numbers, I knew that the solution set is discrete.

I also knew that my graph would occur only in the first quadrant.

I isolated y so I could enter the inequality into my graphing calculator.

I adjusted the calculator window to show only the first quadrant, since the domain and range are both the set of whole numbers. The boundary is a dashed line, which means that the solution set does not include values on the line.

I used the test point $(0, 0)$ to verify that the correct half-plane was shaded. Since $(0, 0)$ is not a solution to the linear inequality, I knew that the half-plane that did not include this point should be shaded. This was done correctly.

When I interpreted the graph, I considered the context of the problem. I knew that

- only discrete points with whole-number coordinates in the solution region made sense.
- points along the dashed boundary are not part of the solution region.
- points with whole-number coordinates along the x -axis and y -axis boundaries are part of the solution region.

I picked two points in the solution region, $(4, 4)$ and $(5, 3)$, as possible solutions to the problem. I verified that each point is a solution to the linear inequality.

Some points in the solution region are more reasonable than others. For example, the point $(1000, 1000)$ is a valid solution, but it might be an unrealistic sales goal.

Sales of four pairs of skis and four snowboards or sales of five pairs of skis and three snowboards will exceed the manager's net revenue goal of more than \$600 a day.

6.1 Page 9

Your Turn

- Would raising the daily sales goal to at least \$1000 change the graph that models this situation? Explain.
- State two combinations of ski and snowboard sales that would meet or exceed this new daily sales goal.

Answers

a)

b)

6.1 Page 10

In Summary p. 220

Key Idea

- When a linear inequality in two variables is represented graphically, its boundary divides the Cartesian plane into two half planes. One of these half planes represents the solution set of the linear inequality, which may or may not include points on the boundary itself.

Need to Know

- To graph a linear inequality in two variables, follow these steps:
 - Step 1.** Graph the boundary of the solution region.
 - If the linear inequality includes the possibility of equality (\leq or \geq), and the solution set is continuous, draw a solid green line to show that all points on the boundary are included.
 - If the linear inequality includes the possibility of equality (\leq or \geq), and the solution set is discrete, stipple the boundary with green points.
 - If the linear inequality excludes the possibility of equality ($<$ or $>$), draw a dashed line to show that the points on the boundary are not included.
 - Use a dashed green line for continuous solution sets.
 - Use a dashed orange line for discrete solution sets.
 - Step 2.** Choose a test point that is on one side of the boundary.
 - Substitute the coordinates of the test point into the linear inequality.
 - If possible, use the origin, (0, 0), to simplify your calculations.
 - If the test point is a solution to the linear inequality, shade the half plane that contains this point. Otherwise, shade the other half plane.
 - Use green shading for continuous solution sets.
 - Use orange shading with green stippling for discrete solution sets.

For example,

$\{(x, y) \mid y \leq -2x + 5, x \in \mathbb{R}, y \in \mathbb{R}\}$

$\{(x, y) \mid y > -2x + 5, x \in \mathbb{R}, y \in \mathbb{R}\}$

$\{(x, y) \mid y \geq -2x + 5, x \in \mathbb{I}, y \in \mathbb{I}\}$

$\{(x, y) \mid y < -2x + 5, x \in \mathbb{I}, y \in \mathbb{I}\}$

- When interpreting the solution region for a linear inequality, consider the restrictions on the domain and range of the variables.
 - If the solution set is continuous, all the points in the solution region are in the solution set.
 - If the solution set is discrete, only specific points in the solution region are in the solution set. This is represented graphically by stippling.
 - Some solution sets may be restricted to specific quadrants. For example, most linear inequalities representing real-world problem situations have graphs that are restricted to the first quadrant.

HOMEWORK...

p. 221: #2, 5, 7, 8, 9

- 1) Declare variables
- 2) State restrictions
- 3) Develop inequation
- 4) Graph solution set

6.1 Page 11

Oct 9-12:36 PM

5.3

Graphing to Solve Systems of Linear Inequalities

GOAL
Solve problems by modelling system

EXPLORE...

- What conclusions can you make about the inequalities graphed below?

system of linear inequalities

A set of two or more linear inequalities that are graphed on the same coordinate plane; the intersection of their solution regions represents the solution set for the system.

SAMPLE ANSWER

Any or all of the following solutions are acceptable:

- It represents a system of two linear inequalities, each with a straight boundary and a solution region.
- One linear inequality is $y \leq -2x + 3$, and the horizontal inequality is $y \geq -2$. I determined $y \leq -2x + 3$ using the slope and y-intercept and the form $y = mx + b$, and I was able to identify $y \geq -2$ because it's a horizontal line through -2 on the y-axis.
- Both inequalities include the possibility of equality because the boundaries are solid.
- The solution set of the system is represented by the overlapping region because it's where the solution regions for the two linear inequalities overlap. The solution set includes points along the boundaries of the overlap.
- The domain and range are from the set of real numbers because the solution region is green and not stippled.
- All four quadrants are included so there are no restrictions on the set of real numbers.

6.3 Page 1

Solving Systems of Linear Inec

A **system of linear inequalities** is an extension of a system of linear equations and consists of two (or more) linear inequalities that have the same variables. For example, $2x + 3y < 4$ and $3x + 4y < 5$ constitute a system of inequalities if x represents the same item in both equations, y represents the same item in both equations, and both equations describe the same context.

Example #1:
Graph the following system and determine a possible solution

$\{(x, y) \mid y \leq 3x - 1, x \in \mathbb{R}, y \in \mathbb{R}\}$

Test (0,0)

$y = 3x - 1$

$0 \leq 3(0) - 1$


$0 \leq -1$ False

$\{(x, y) \mid y > -\frac{1}{3}x + 4, x \in \mathbb{R}, y \in \mathbb{R}\}$


Test (0,0)

$0 > 4$ False

Oct 15-10:57 AM



Feb 5-2:21 PM



Sep 12-9:02 AM

APPLY the Math Can be found on p.230

EXAMPLE 2 Solving graphically a system of two linear inequalities with continuous variables

Graph the solution set for the following system of inequalities. Choose two possible solutions from the set.

$3x + 2y > -6$
 $y \leq 3$

Peter's Solution: Using graph paper

$x \in \mathbb{R}, y \in \mathbb{R}$

$3x + 2y > -6$
 x-intercept: $3x + 2(0) = -6 \Rightarrow x = -2$
 y-intercept: $3(0) + 2y = -6 \Rightarrow y = -3$
 Points: $(-2, 0)$ and $(0, -3)$


Test $(0, 0)$ in $3x + 2y > -6$.
 $3(0) + 2(0) = 0 > -6$. Since $0 > -6$, $(0, 0)$ is in the solution region.

$y \leq 3$

The overlapping solution region represents the solution set of the system of linear inequalities. Therefore, $(2, -3)$ and $(-1.5, 3)$ are two possible solutions.

Your Turn
 How would the solution region change if $x \in \mathbb{I}$ and $y \in \mathbb{I}$?
 How would it stay the same?

6.3 Page 6



Feb 5-2:41 PM

Let's check out the web...

EXAMPLE #3: Applet from online Math Tutor

"Khan"

Oct 15-11:05 AM

In Summary

Key Ideas

- When graphing a system of linear inequalities, the boundaries of its solution region may or may not be included, depending on the types of linear inequalities (\geq , \leq , $<$, or $>$) in the system.
- Most systems of linear inequalities representing real-world situations are restricted to the first quadrant because the values of the variables in the system must be positive.

Need to Know

- Any point in the solution region for a system is a valid solution, but some solutions may make more sense than others depending on the context of the problem.
- You can validate a possible solution from the solution region by checking to see if it satisfies each linear inequality in the system. For example, to validate if $(2, 2)$ is a solution to the system:

$$\begin{aligned} x + y &\geq 1 \\ 2 > x - 2y \end{aligned}$$

Validating $(2, 2)$ for $x + y \geq 1$:		Validating $(2, 2)$ for $2 > x - 2y$:	
LS	RS	LS	RS
$x + y$	1	2	$x - 2y$
$2 + 2$			$2 - 2(2)$
4			-2
$4 \geq 1$	valid	$2 > -2$	valid
- Use an open dot to show that an intersection point of a system's boundaries is excluded from the solution set. An intersection point is excluded when a dashed line intersects either a dashed or solid line.
- Use a solid dot to show that an intersection point of a system's boundaries is included in the solution set. This occurs when both boundary lines are solid.

6.3 Page 11

HOMWORK...

Puzzle Worksheet - Systems of Linear Inequations.docx

Worksheet - Systems of Linear Inequations.docx

page 225 #1,2

Sep 12-9:56 AM

SOLUTIONS...

PUZZLE WORKSHEET:

What Did the Toothless Old Termite Say When He Entered a Tavern?

Graph each system of inequalities. Shade and describe the solution set of the system using a word or phrase. Write the coordinates of any vertices of the solution set in the space provided.

1) $x + y \leq 1$
 $x - y \geq -3$

2) $x + 2y \leq 12$
 $x - y \leq 1$

3) $x + y \leq 1$
 $x - y \geq -3$

4) $x + y \leq 1$
 $x - y \geq -3$

5) $4x + y < 2$
 $y > -2$

6) $3x + 2y \geq -2$
 $x + 2y \leq 2$

WORKSHEET:

1) $y \leq -x - 2$
 $y \geq -5x + 2$

2) $y > -x - 2$
 $y < -5x + 2$

3) $y \leq \frac{1}{2}x + 2$
 $y < -2x - 3$

4) $x \leq -3$
 $y < \frac{3}{5}x + 2$

5) $4x + y < 2$
 $y > -2$

6) $3x + 2y \geq -2$
 $x + 2y \leq 2$

Sep 12-10:11 AM

WARM-UP: Graph the solution and state 2 possible solutions...

$\{(x,y) | 2x + y > 8, x \in W, y \in W\}$
 $\{(x,y) | y \leq 2, x \in W, y \in W\}$

$y > -2x + 8$
 $y \leq 2$

$(6,0)$
 $(10,-4)$

Oct 16-10:42 AM

Applications: Systems Involving Inequalities

STEP 1 - Declare Variables
State Restrictions

STEP 2 - Create Linear Inequalities

STEP 3 - Graph Solution Set

STEP 4 - Answer question(s)

EXAMPLE #1:
To raise funds for π -day, the PI Committee has 500 T-shirts to sell. \leftarrow
They have two varieties:
#1. 'I 8 Sum π ' or #2. ' π -DAY 2013'.
They expect to sell at least twice as many of the first as the second. \leftarrow

a) Define the variables and restrictions. Write a system of linear inequalities that models the situation.

$x = \pi$ -shirt, $y = \pi$ -DAY
 $x \geq 2y$ \leftarrow hand bit
 $x + y \leq 500$

b) Graph the system of inequalities.

c) State a combination of T-shirt sales.

200 1
 100 2

Oct 16-10:46 AM

5.2 Exploring Graphs of Systems of Linear Inequalities

GOAL
Explore graphs of situations that can be modelled by systems of two linear inequalities in two variables.

EXPLORE the Math

A nursery school serves morning and afternoon snacks to its students. The morning snacks are fruits, vegetables, and juice, and the afternoon snacks are cheese and milk.

- The school can accommodate 50 students or fewer altogether. Students can attend for just the morning or for a full day.
- The morning snack costs \$1 per student per week, and the afternoon snack costs \$2 per student per week.
- The weekly snack budget is \$120 or less.

? What combinations of morning and full-day students can the school accommodate and stay within the weekly snack budget?

? What combinations of morning and full-day students can the school accommodate and stay within the weekly snack budget?

Sample Solution

First, we represented the two unknowns in the problem using x and y :

- x is the number of morning students.
- y is the number of full-day students.

Then we wrote a linear inequality to represent each part of the problem:

- The total cost of the snacks, as it relates to the number of students, is the sum of the cost of the morning snack multiplied by the number of morning students and the sum of the cost of the afternoon snack multiplied by the number of full-day students. The total cost is \$120 or less:
 $x + 3y \leq 120$ \leftarrow cost
- The total number of students can be up to and including 50 students:
 $x + y \leq 50$

Finally, we graphed both linear inequalities on the same coordinate plane.

$x + 3y \leq 120 \rightarrow y \leq \frac{1}{3}x + 40$
 $x + y \leq 50 \rightarrow y \leq -x + 50$

Full-day Students and Morning Students


For the exploration, there is no independent-variable relationship between the two variables. Therefore, students can represent either type of student (morning or full day) along the x -axis. As a result, graphical representations may appear different but yield the same solution. Some students may use guess and check to find several solutions that work for both inequalities. Other students may realize that they can graph each inequality on the same coordinate plane and look for the intersection of the solution region.

We knew that the area where the two solution regions intersect or overlap contains points that represent all the possible combinations of morning and full-day students that will work. We also knew that only whole-number points, such as (24, 24) and (8, 36), make sense.

In Summary


Key Ideas

- Some contextual situations can be modelled by a system of two or more linear inequalities.
- All of the inequalities in a system of linear inequalities are graphed on the same coordinate plane. The region where their solution regions intersect or overlap represents the solution set to the system. For example, this graph shows the solution region to this system:
 $\{(x, y) | y \geq x, x \in \mathbb{R}, y \in \mathbb{R}\}$
 $\{(x, y) | y \leq 5, x \in \mathbb{R}, y \in \mathbb{R}\}$



Need to Know

- As with the solution region for a single linear inequality, the solution region for a system of linear inequalities can be discrete or continuous and can be restricted to certain quadrants. For example, the graph to the right shows the system described below:
 $\{(x, y) | y \geq 1, x \in \mathbb{W}, y \in \mathbb{W}\}$
 $\{(x, y) | y \leq -3x + 6, x \in \mathbb{W}, y \in \mathbb{W}\}$
 Its solution region is restricted to discrete points with whole-number coordinates in the first quadrant.
- If the solution regions for the linear inequalities in the system do not overlap, there is no solution.



6.2 Page 5

Logic - Hockey Time.doc

INSTRUCTIONS:
 Complete the grid below by using a '✓' for facts and 'X' for elimination

	Newcastle	Renous	Tabusintac	Douglasstown	Millerton	Cul Tech	Eng	OP	Bio	Math	Van	Tor	Mtl	Edm	Ott
March 26	X	X	X	X	X										
May 3		X	X												
May 14		X	X												
November 13		X	X												
November 21		X	X												
Van															
Tor															
Mtl															
Edm															
Ott															
Cul Tech															
Eng															
OP															
Bio															
Math															

Sep 13-8:51 AM

HOMWORK...

Logic - Hockey Time.doc
 (Due on Friday FIRST of class)

p. ~~225: #1 & 2~~

p. 235: #2, 5 & 6

QUIZ on Friday!!!


Oct 15-11:16 AM

Logic - Hockey Time.doc

INSTRUCTIONS:
 Complete the grid below by using a '✓' for facts and 'X' for elimination

	Newcastle	Renous	Tabusintac	Douglasstown	Millerton	Cul Tech	Eng	OP	Bio	Math	Van	Tor	Mtl	Edm	Ott
March 26	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
May 3	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
May 14	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
November 13	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
November 21	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Van	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Tor	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Mtl	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Edm	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Ott	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Cul Tech	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Eng	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
OP	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Bio	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Math	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

Sep 13-8:51 AM



Feb 7-12:46 PM

LEARN ABOUT the Math *** Can be found on p. 226

A computer makes two types of boats, an aluminum fishing boat and a fiberglass fishing boat. The aluminum fishing boats are more expensive than the fiberglass fishing boats. The demand for fiberglass boats is greater than the demand for aluminum boats, so the company makes at least 5 more fiberglass boats than aluminum boats each day.

What combination of boats should the company make each day?

Volume 1 Solving a problem with discrete whole-number variables using a system of inequalities

Mary's Solution: Using graph paper

Let x represent the number of aluminum fishing boats. Let y represent the number of fiberglass fishing boats. $x \in \mathbb{W}$ and $y \in \mathbb{W}$.

The relationship between the two types of boats can be represented by the system of inequalities: $x + y \leq 20$ and $x \geq 5 + y$.

Step 1: Graph each linear inequality. I know that to graph an inequality on a coordinate plane, I need to graph the boundary line and shade the correct half-plane. To graph each boundary, I write each inequality as an equation and then determine the x - and y -intercepts. I could also use a graphing calculator.

Step 2: Test a point in each half-plane. I know that a point is a solution to a system of inequalities if it satisfies both inequalities. I test the point $(10, 10)$ in the system of inequalities. $10 + 10 \leq 20$ is true, but $10 \geq 5 + 10$ is false. So $(10, 10)$ is not a solution to the system of inequalities. I test the point $(5, 15)$ in the system of inequalities. $5 + 15 \leq 20$ is true, and $5 \geq 5 + 15$ is false. So $(5, 15)$ is not a solution to the system of inequalities. I test the point $(15, 5)$ in the system of inequalities. $15 + 5 \leq 20$ is true, and $15 \geq 5 + 5$ is true. So $(15, 5)$ is a solution to the system of inequalities. I shade the region that contains $(15, 5)$.

Step 3: Determine the solution region. The solution region is the shaded region in the first quadrant. The solution region is discrete because the number of boats is a whole number.

Fiberglass vs. Aluminum


Number of aluminum fishing boats (x) vs. Number of fiberglass fishing boats (y).

Step 4: Graph the solution set for the system. I use the same coordinate plane. I use the points $(15, 5)$ and $(10, 10)$ to draw a grid. I shade the region that contains $(15, 5)$ and $(10, 10)$.


Step 5: Test a point in each half-plane. I know that a point is a solution to a system of inequalities if it satisfies both inequalities. I test the point $(15, 5)$ in the system of inequalities. $15 + 5 \leq 20$ is true, and $15 \geq 5 + 5$ is true. So $(15, 5)$ is a solution to the system of inequalities. I shade the region that contains $(15, 5)$.

Step 6: Determine the solution region. The solution region is the shaded region in the first quadrant. The solution region is discrete because the number of boats is a whole number.

6.3 Page 2



Feb 7-2:28 PM



Feb 7-1:06 PM

Reflecting

- A. Is every point on the boundaries of the solution region a possible solution? Explain.
- B. Are the three points where the boundaries intersect part of the solution region? Explain.
- C. How would the graph change if fewer than 25 boats were made each day?
- D. All points with whole-number coordinates in the solution region are valid, but are they all reasonable? Explain.

Answers

A.
 B.
 C.
 D.

6.3 Page 5

$x \rightarrow$ younger $x \leq w$
 $y \rightarrow$ older $y \leq w$

$$x + y \leq 36$$

$$x \geq 2y$$

$$\frac{-2y}{-2} \geq \frac{-x}{-2}$$

$$y \leq \frac{1}{2}x$$

Feb 7-3:01 PM

7241

6. $l < 100$ $l > 0, l \in \mathbb{R}$
 $w > 0, w \in \mathbb{R}$
 $2l + 2w \leq 400$

7. $P + M \leq 30$
 $18P + 10M \leq 470$

Feb 7-3:16 PM

*** Can be found on p. 233

EXAMPLE 3 Solving graphically a problem with continuous positive variables

A hoop is a sailboat with two sails: a mainsail and a jib. When a sail is fully out or up, it is said to be "out 100%." When the winds are high, sailors often reef, or pull in, the sails to be less than their full capability.

- Jim is sailing in winds of 22 knots, so he wants no more than 80% of the mainsail out.
- Jim also wants more mainsail out than jib.

What possible combinations of mainsail and jib can Jim have out?

Louise's Solution: Using graph paper

Let m represent the percent of mainsail out.
 Let j represent the percent of jib out.

$m \geq 0$ and $j \geq 0$, where $m, j \in \mathbb{R}$

The relationship between the two types of sails can be represented by the following system of two linear inequalities:

$m \leq 80$
 $j < m$

$m \leq 80$
 Boundary: $m = 80$
 Boundary is a vertical line with an m -intercept of 80.

$j < m$
 Boundary: $j = m$
 Boundary line has a slope of 1 and a j -intercept of 0.

The inequalities describe the following information:

- No more than 80% of the mainsail can be out.
- Less jib than mainsail must be out.

I decided to use m as the independent variable. I examined each inequality to determine its boundary.

- Since m is the independent variable, I knew the boundary for $m \leq 80$ would be a vertical line through $m = 80$.
- I knew the boundary of $j < m$ has a slope of 1 and passes through the point $(0, 0)$.

For $m = 80$, I drew a solid green vertical line through $m = 80$ and then shaded the half plane to its left green, since the inequality sign is \leq .

For $j < m$, I drew a green dashed line through $(0, 0)$ with a slope of 1 and I shaded the half plane below green since the inequality is $<$.

I drew an open dot where the dashed boundary intersects the solid boundary to show that point isn't part of the solution region.

The solution region for the system is a right triangle and consists of all the points in the overlapping region, including the solid boundary and the m -axis from 0 to 80.

I looked for several solutions in the solution region. I knew that I could choose points with decimal coordinates since the solution region is continuous.

Percent jib vs. Percent Mainsail

Percent of jib out vs. Percent of mainsail out

Percent of jib out vs. Percent of mainsail out

$\{(m, j) \mid m \leq 80, m \geq 0, j \geq 0, m \in \mathbb{R}, j \in \mathbb{R}\}$
 $\{(m, j) \mid j < m, m \geq 0, j \geq 0, m \in \mathbb{R}, j \in \mathbb{R}\}$

Any point in the solution region represents an acceptable combination. For example,

- 80% of the mainsail and 70% of the jib can be out.
- 70% of the mainsail and 40% of the jib can be out.
- 40.5% of the mainsail and 30.7% of the jib can be out.

6.3 Page 9

HOMEWORK...

p. 236: #7 - 10

NOTE: Each question requires a graph to get possible solutions!

Quiz on graphing inequalities on **TOMORROW!!!**

5.4 Optimization Problems I: Creating the Model

optimization problem

A problem where a quantity must be maximized or minimized following a set of guidelines or conditions.

objective function

In an optimization problem, the equation that represents the relationship between the two variables in the system of linear inequalities and the quantity to be optimized.

feasible region

The solution region for a system of linear inequalities that is modelling an optimization problem.

constraint

A limiting condition of the optimization problem being modelled, represented by a linear inequality.

Need to Know

- You can create a model for an optimization problem by following these steps:
 - Step 1.** Identify the quantity that must be optimized. Look for key words, such as *maximize* or *minimize*, *largest* or *smallest*, and *greatest* or *least*.
 - Step 2.** Define the variables that affect the quantity to be optimized. Identify any restrictions on these variables.
 - Step 3.** Write a system of linear inequalities to describe all the constraints of the problem. Graph the system.
 - Step 4.** Write an objective function to represent the relationship between the variables and the quantity to be optimized.

Mar 15-1:32 PM

6.4 Page 1

APPLY the Math

EXAMPLE 1 Creating a model for an optimization problem with whole-number variables

Three teams are travelling to a basketball tournament in cars and minivans.

- Each team has no more than 2 coaches and 14 athletes.
- Each car can take 4 team members, and each minivan can take 6 team members.
- No more than 4 minivans and 12 cars are available.

The school wants to know the combination of cars and minivans that will require the **minimum** and **maximum** number of vehicles. Create a model to represent this situation.

Juanita's Solution

Let m represent the number of minivans.
Let c represent the number of cars.

The two variables in the problem are the number of cars and the number of minivans. The values of these variables are whole numbers.

$m \in \mathbb{W}$ and $c \in \mathbb{W}$

Constraints:

- Number of cars available: $c \leq 12$
- Number of minivans available: $m \leq 4$
- Number of team members: $4c + 6m \leq 48$

I knew that this is an **optimization problem** because the number of vehicles has to be minimized and maximized.

I wrote three linear inequalities to represent the three limiting conditions, or **constraints**.

The maximum number of team members is the number of teams multiplied by the maximum number of coaches and athletes: $3(14) + 3(2) = 48$

Objective function:
Let V represent the total number of vehicles.

$V = c + m$

I created an equation, called the **objective function**, to represent the relationship between the two variables (number of minivans and number of cars) and the quantity to be minimized and maximized (number of vehicles).

I graphed the system of three inequalities.

One of the solutions in the **feasible region** represents the combination of cars and minivans that results in the minimum total number of vehicles and another solution represents the maximum. I think I could use the objective function to determine each point, but I am not certain how yet.

feasible region
The solution region for a system of linear inequalities that is modelling an optimization problem.

The graph shows the feasible region for the optimization problem. The x-axis is 'Number of minivans' (0 to 10) and the y-axis is 'Number of cars' (0 to 14). The constraints are: $c \leq 12$, $m \leq 4$, and $4c + 6m \leq 48$. The feasible region is shaded in green. Handwritten notes include: $m=4$, $c=6$, $V=m+c$, $V=4+6$, $V=10$. Other notes include $16 \times 3 = 48$ and $m=4$, $c=6$.

6.4 Page 4

Number of Cars and Minivans

The graph shows the feasible region for the optimization problem. The x-axis is 'Number of minivans' (0 to 10) and the y-axis is 'Number of cars' (0 to 14). The constraints are: $c \leq 12$, $m \leq 4$, and $4c + 6m \leq 48$. The feasible region is shaded in green. Handwritten notes include: $m=4$, $c=6$, $V=m+c$, $V=4+6$, $V=10$. Other notes include $16 \times 3 = 48$ and $m=4$, $c=6$.

Feb 10-1:12 PM

Your Turn

Suppose that the greatest number of athletes changed from 14 to 12 per team. How would Juanita's model change?

Answer

In the original model, the greatest number of athletes was 14. If this number changes to 12, the model must be adjusted to reflect this new constraint. The feasible region would become smaller, and the maximum value of the objective function would likely decrease.


6.4 Page 5

EXAMPLE 2 Creating a model for a maximization problem with positive real-number variables

A refinery produces oil and gas.

- At least 2 L of gasoline is produced for each litre of heating oil.
- The refinery can produce up to 9 million litres of heating oil and 6 million litres of gasoline each day.
- Gasoline is projected to sell for \$1.10 per litre. Heating oil is projected to sell for \$1.75 per litre.

The company needs to determine the daily combination of gas and heating oil that must be produced to maximize revenue. Create a model to represent this situation.



Umberto's Solution

Let h represent the number of litres of heating oil. Let g represent the number of litres of gasoline.

Restrictions:
 $h \geq 0$ and $g \geq 0$, where $h \in \mathbb{R}$ and $g \in \mathbb{R}$


Constraints:
 Ratio of gasoline produced to oil produced:
 $g \geq 2h$
 Amount of gasoline that can be produced:
 $g \leq 6\,000\,000$
 Amount of oil that can be produced:
 $h \leq 9\,000\,000$

Let R represent total revenue from sales of gasoline and heating oil.
 Objective function to maximize:
 $R = 1.10g + 1.75h$

Handwritten notes: "I knew that this is an optimization problem because the total revenue has to be maximized." "The two variables in the problem are the volume of heating oil and the volume of gasoline, both in litres. Litres are measured using positive real numbers." "I created inequalities to represent the five constraints of the problem." "I treated the restrictions on each variable as a constraint." "I wrote an objective function to represent the relationship between the two variables (volume of heating oil and volume of gasoline) and the quantity to be maximized (total revenue)." "I graphed the system of inequalities in the 1st quadrant because of the restrictions on the variables. The feasible region is a right triangle and includes all points on its boundaries." "I think I can use the objective function to determine which point in the feasible region represents the combination of oil and gas that will result in the maximum revenue, but I am not sure how yet."

Handwritten note: "don't graph"

Production of Gasoline and Heating Oil



6.4 Page 6

In Summary


Key Ideas

- To solve an optimization problem, you need to determine which combination of values of two variables results in a maximum or minimum value of a related quantity.
- When creating a model, the first step is to represent the situation algebraically. An algebraic model includes these parts:
 - a defining statement of the variables used in your model
 - a statement describing the restrictions on the variables
 - a system of linear inequalities that describes the constraints
 - an objective function that shows how the variables are related to the quantity to be optimized
- The second step is to represent the system of linear inequalities graphically.
- In optimization problems, any restrictions on the variables are considered constraints. For example, if you are working with positive real numbers, $x \geq 0$ and $y \geq 0$ are constraints and should be included in the system of linear inequalities.

Need to Know

- You can create a model for an optimization problem by following these steps:
 - Step 1.** Identify the quantity that must be optimized. Look for key words, such as *maximize* or *minimize*, *largest* or *smallest*, and *greatest* or *least*.
 - Step 2.** Define the variables that affect the quantity to be optimized. Identify any restrictions on these variables.
 - Step 3.** Write a system of linear inequalities to describe all the constraints of the problem. Graph the system.
 - Step 4.** Write an objective function to represent the relationship between the variables and the quantity to be optimized.

Mar 19-9:38 PM



Feb 10-1:24 PM

HOMEWORK...

Page 248: #1abc, #2, #3

Stape "hockey"

NOTE:

Create a model means graph the solution region

Quiz → Tuesday

QUIZ TIME...

When finished pass your quiz in and work on the following:

GOAL
Solve optimization problems.

EXPLORE...

The following system of linear inequalities has been graphed below:

System of linear inequalities:
 $y \geq 0$
 $x \geq 0$
 $y \leq 4$
 $x \leq 3$
 $3y \geq -x + 9$

$y = -\frac{1}{3}x + 3$

$T = 5x + y$

$T = 5(0) + 4 = 4$
 $T = 5(0) + 3 = 3$
 $T = 5(3) + 2 = 17$
 $T = 5(0) + 0 = 0$

a) For each objective function, what points in the feasible region represent the minimum and maximum values?
 i) $T = 5x + y$
 ii) $T = x + 5y$

b) What do you notice about the optimal points for the two objective functions? Why do you think this happened?

SAMPLE ANSWER

a) i) For $T = 5x + y$,

If (x, y) is...	Then...	
(3, 2)	$T = 5(3) + 2 = 17$	
(3, 4)	$T = 5(3) + 4 = 19$	maximum
(0, 3)	$T = 5(0) + 3 = 3$	minimum
(0, 4)	$T = 5(0) + 4 = 4$	

ii) For $T = x + 5y$,

If (x, y) is...	Then...	
(3, 2)	$T = 3 + 5(2) = 13$	minimum
(3, 4)	$T = 3 + 5(4) = 23$	maximum
(0, 3)	$T = 0 + 5(3) = 15$	
(0, 4)	$T = 0 + 5(4) = 20$	

b) I noticed that the values of the coefficients of the variables and the values of the variables themselves all contribute to the value of the objective function. For $T = 5x + y$, the x-value is multiplied by 5 and the y-value is multiplied by 1. For $T = x + 5y$, the x-value is multiplied by 1 and the y-value is multiplied by 5. In each case, the greater the coordinate that is multiplied by 5, the greater the value of the objective function is. The converse is true for the least values.

EXAMPLE of an OPTIMIZATION Problem...

Mick and Keith make MP3 covers to sell, using beads and stickers.

- At most, 45 covers with stickers and 55 bead covers can be made per day.
- Mick and Keith can make 45 or more covers, in total, each day.
- It costs \$0.75 to make a cover with stickers, \$1.00 to make one with beads.



Let x represent the number of covers with stickers and let y represent the number of bead covers. Let C represent the cost of making the covers.

RESTRICTIONS:

CONSTRAINTS:


OBJECTIVE FUNCTION:

a) Graph the solution set.

b) What are the vertices of the feasible region?

c) Which point would result in the maximum value of the objective function?

d) Which point would result in the minimum value of the objective function?



HOMEWORK...

p. 252: #1 - 3

p. 248: #4 - 6

5.5 Optimization Problems II: Exploring Solutions

Need to Know

- The solution to an optimization problem is usually found at one of the vertices of the feasible region.
- To determine the optimal solution to an optimization problem using linear programming, follow these steps:
 - Step 1.** Create an algebraic model that includes:
 - a defining statement of the variables used in your model
 - the restrictions on the variables
 - a system of linear inequalities that describes the constraints
 - an objective function that shows how the variables are related to the quantity to be optimized
 - Step 2.** Graph the system of inequalities to determine the coordinates of the vertices of its feasible region.
 - Step 3.** Evaluate the objective function by substituting the values of the coordinates of each vertex.
 - Step 4.** Compare the results and choose the desired solution.
 - Step 5.** Verify that the solution(s) satisfies the constraints of the problem situation.

optimal solution
A point in the solution set that represents the maximum or minimum value of the objective function.

Mar 19-9:53 PM

GOAL
Explore the feasible region of a system of linear inequalities.

EXPLORE the Math

A toy company manufactures two types of toy vehicles: racing cars and sport-utility vehicles.

- Because the supply of materials is limited, no more than 40 racing cars and 60 sport-utility vehicles can be made each day.
- However, the company can make 70 or more vehicles, in total, each day.
- It costs \$8 to make a racing car and \$12 to make a sport-utility vehicle.

There are many possible combinations of racing cars and sport-utility vehicles that could be made. The company wants to know what combinations will result in the minimum and maximum costs, and what those costs will be.

The following model represents this situation. The feasible region of the graph represents all the possible combinations of racing cars (r) and sport-utility vehicles (s).

Variables:
Let s represent the number of sport-utility vehicles.
Let r represent the number of racing cars.
Let C represent the cost of production.

Restrictions:
 $s \in \mathbb{W}, r \in \mathbb{W}$

Constraints:
 $s \geq 0$
 $r \geq 0$
 $r \leq 40$
 $s \leq 60$
 $r + s \geq 70$

Objective function to optimize:
 $C = 12s + 8r$

? How can you use patterns in the feasible region to predict the combinations of sport-utility vehicles and racing cars that will result in the minimum and maximum values of the objective function?

Value of $C = 8r + 12s$

- as s increases: 720 at (40, 30), 840 at (50, 30), 960 at (60, 30)
- as r increases: 860 at (45,40), 780 at (45, 30) 740 at (45, 25)
- in the middle of the solution region: 820 at (45, 35), 880 at (50, 35), 840 at (50, 30)
- at the corners of the solution region: 800 at (60, 10), 1040 at (60, 40), 680 at (30, 40)

6.5 Page 1

Reflecting

- With a partner, discuss the pattern in the value of C throughout the feasible region. Is the pattern what you expected? Explain.
- As you move from left to right across the feasible region, what happens to the value of C ?
- As you move from the bottom to the top of the feasible region, what happens to the value of C ?
- What points in the feasible region result in each **optimal solution**?
 - the maximum possible value of C
 - the minimum possible value of C
- Explain how you could verify that your solutions from part D satisfy each constraint in the model.

Answers

-
-
-
- -
- -

6.5 Page 3

EXAMPLE #1...

The vertices of the feasible region of a graph of a system of linear inequalities are $(-4, -8)$; $(5, 0)$ and $(1, -6)$. Which point would result in the minimum value of the objective function $C = 0.50x + 0.60y$?

$C = 0.50(-4) + 0.60(-8) = -6.8$
 $C = 0.50(5) + 0.60(0) = 2.5$
 $C = 0.50(1) + 0.60(-6) = -3.1$

Oct 22-11:02 AM

EXAMPLE #2...
 The following model represents an optimization problem. Determine the maximum solution.
 Restrictions: $x \in \mathbb{R}$ and $y \in \mathbb{R}$
 Constraints: $y \leq 1$; $2y \geq -3x + 2$; $y \geq 3x - 8$
 Objective Function: $D = -4x + 3y$

Oct 22-11:13 AM

EXAMPLE #2...
 The following model represents an optimization problem. Determine the maximum solution.
 Restrictions: $x \in \mathbb{R}$ and $y \in \mathbb{R}$
 Constraints: $y \leq 1$; $2y \geq -3x + 2$; $y \geq 3x - 8$
 Objective Function: $D = -4x + 3y$

$(0,1) \rightarrow D = -4(0) + 3(1) = 3$
 $(3,1) \rightarrow D = -4(3) + 3(1) = -9$
 $(2,-2) \rightarrow D = -4(2) + 3(-2) = -14$

Oct 22-11:13 AM

Four MVHS teams are travelling to a basketball tournament in cars and minivans.

- Each team has no more than 2 coaches and 10 athletes
- Each car can take 4 team members. Each minivan can take 6 team members.
- No more than 6 cars are available, but more than 3 minivans are available.

Mr. Watters wants to know the combination of cars and minivans that will require the maximum number of vehicles...

a) Create an algebraic model to represent this situation.

b) Graph the model.

$4x + 6y \leq 48$
 $x \leq 6$
 $y > 3$
 $V = x + y$

c) What combination of cars/minivans will result in the maximum number of vehicles?

d) How many team members can travel in the maximum number of vehicles?

$4x + 6y = 48$
 x -int y -int
 $4x + 6(0) = 48$ $4(6) + 6y = 48$
 $4x = 48$ $6y = 48$
 $x = 12$ $y = 8$
 $(12, 0)$ $(0, 8)$
 $V = x + y$
 $(0, 8) \rightarrow V = 8$
 $(6, 4) \rightarrow V = 10$
 $(0, 3)$
 $(6, 3?)$

Oct 22-11:17 AM

In Summary

Key Ideas

- The value of the objective function for a system of linear inequalities varies throughout the feasible region, but in a predictable way.
- The optimal solutions to the objective function are represented by points at the intersections of the boundaries of the feasible region. If one or more of the intersecting boundaries is not part of the solution set, the optimal solution will be nearby.

Need to Know

- You can verify each optimal solution to make sure it satisfies each constraint by substituting the values of its coordinates into each linear inequality in the system.
- The intersection points of the boundaries are called the vertices, or corners, of the feasible region.

HOMEWORK...

- Page 259: #1-4

WARM UP - use graph paper

e.g., *Problem:* A library is buying both hardcover and paperback books. It plans to purchase at most four times as many paperbacks as hardcover books. Altogether the plan is to purchase no fewer than 200 books. Hardcover books average \$35.75 in cost while paperbacks average \$12.20. How can the library minimize its costs?

Solution: Let x represent the number of hardcover books. Let y represent the number of paperback books. Let C represent the total cost of the books.

Objective function to minimize: $C = 35.75x + 12.2y$

Constraints and restrictions:

$$\{(x, y) \mid x + y \geq 200, x \in \mathbb{W}, y \in \mathbb{W}\}$$

$$\{(x, y) \mid 4x \geq y, x \in \mathbb{W}, y \in \mathbb{W}\}$$

The library should purchase 40 hardcover books and 160 paperback books, for a total cost of \$3382.00.

Mar 19-9:46 PM

Oct 23-8:15 AM

Restrictions

Constraints

ObjectiveFunction

e.g., *Problem:* A library is buying both hardcover and paperback books. It plans to purchase at most four times as many paperbacks as hardcover books. Altogether the plan is to purchase no fewer than 200 books. Hardcover books average \$35.75 in cost while paperbacks average \$12.20. How can the library minimize its costs?

Oct 23-9:36 AM

ONE MORE...

Malia and Lainey are baking cupcakes and banana mini-loaves to sell at a school fundraiser...

- No more than 60 cupcakes and 35 mini-loaves can be made each day.
- Malia and Lainey can mke no more than 80 baked goods, in total, each day.
- It costs \$0.50 to make a cupcake and \$0.75 to make a mini-loaf.

Determine the maximum cost to produce the baked goods.

Variables & restrictions
 $c \rightarrow$ cupcakes $c \in \mathbb{W}$
 $m \rightarrow$ mini loaf $m \in \mathbb{W}$

constraints
 $c \geq 0$ $c \leq 60$
 $m \geq 0$ $m \leq 35$
 $m + c \leq 80$


ObjectiveFunction
 $C = 0.5c + 0.75m$

Oct 22-11:08 AM

ONE MORE...

Malia and Lainey are baking cupcakes and banana mini-loaves to sell at a school fundraiser...

- No more than 60 cupcakes and 35 mini-loaves can be made each day.
 - Malia and Lainey can mke no more than 80 baked goods, in total, each day.
 - It costs \$0.50 to make a cupcake and \$0.75 to make a mini-loaf.
- Determine the minimum cost to produce the baked goods.



Oct 22-11:08 AM

HOMEWORK...

p. 261: #5, 7, 8, 11, 13

Oct 23-11:50 AM

HOMEWORK: Test is on THURSDAY!!

Review/Practice Questions...

- p. 239: Mid-Chapter Review (Frequently Asked Questions) *
- p. 241: Mid-Chapter Practice Questions
- p. 266: Chapter Review (Frequently Asked Questions) *
- p. 267: Chapter Practice Questions
- p. 265: Chapter Self-Test (Do this AFTER you practice)


LOGIC PROBLEM is due

Tuesday!

Logic - Inequality Sudoku #2.doc

Sep 20-9:23 AM

inequalities.



$y = 2x - 3$
 $y \geq 2x - 3$
 $0 \geq 2(0) - 3$
 $0 \geq -3$
 True

$y = -2x + 7$
 $y \leq -2x + 7$
 $0 < 7$
 True

Determine the linear equation that presents the boundary of each linear inequality. Explain how you determined each equation. Represent the system of linear inequalities graphically.

Feb 19-12:43 PM

p268.
 #7. $m \rightarrow$ male $m \in \mathbb{W}$
 $f \rightarrow$ female $f \in \mathbb{W}$


$m \geq 3f$
 $m + f \leq 28$

graph!
 find vertices

Objective function
 $R = 115m + 90f$

$m \geq 3f$
 $12 = 3(4)$

Max (,)



Sep 20-10:01 AM

Feb 19-2:30 PM

Attachments

NOTES - Graphing Linear Relationships.docx

Puzzle Worksheet - Graphing Lines.docx

Puzzle Worksheet - Graphing Inequalities with One Variable.docx

fm6s1-p5.tns

6Ws1e1.mp4

6Ws1e2.mp4

Worksheet - Graphing Linear Inequalities.pdf

fm6s1-p9.tns

6Ws1e3.mp4

6Ws3e2.mp4

6Ws4e1.mp4

Puzzle Worksheet - Graphing Linear Inequalities with Two Variables.pdf

Worksheet - Solving Systems of Linear Inequalities.pdf

Warm Up - Prior Knowledge for Coordinate Geometry.docx

NOTES - Graphing a Linear Inequation.docx

Logic - Three Little Pigs.doc

Puzzle Worksheet - Systems of Linear Inequations.docx

Worksheet - Systems of Linear Inequations.docx

Example - Application of a Linear Inequality.docx

Logic - Hockey Time.doc

Logic - Inequality Sudoku #2.doc