

Here is a more **Mathematical** definition...

■ ABSOLUTE VALUE

E.1 DEFINITION. The *absolute value* or *magnitude* of a real number a is denoted by $|a|$ and is defined by

$$|a| = \begin{cases} a & \text{if } a \geq 0 \\ -a & \text{if } a < 0 \end{cases}$$

► Example 1

$$|5| = 5 \quad \left| -\frac{4}{7} \right| = -\left(-\frac{4}{7} \right) = \frac{4}{7} \quad |0| = 0 \blacktriangleleft$$

Since $5 > 0$

Since $-\frac{4}{7} < 0$

Since $0 \geq 0$

Note that the effect of taking the absolute value of a number is to strip away the minus sign if the number is negative and to leave the number unchanged if it is nonnegative.

Textbook Pg. 304

$$\#21 \quad t_m = \sqrt{\frac{h}{1.8}} \quad t_E = \sqrt{\frac{h}{4.9}}$$

$$\begin{matrix} \approx 0.2 \\ 0.3 - 0.2 \\ \times \end{matrix} \quad \sqrt{\frac{h}{1.8}} - \sqrt{\frac{h}{4.9}} = 0.5$$

$$\left(\sqrt{\frac{h}{1.8}}\right)^2 = \left(0.5 + \sqrt{\frac{h}{4.9}}\right)^2$$

$$\frac{1}{1.8}h - \frac{1}{4.9}h - 0.25 = \sqrt{\frac{h}{4.9}} + \frac{h}{4.9}$$

$$0.5h - 0.204h - 0.25 = \sqrt{\frac{h}{4.9}}$$
$$(0.352h - 0.25)^2 = \left(\sqrt{\frac{h}{4.9}}\right)^2$$

$$\begin{matrix} \times 4.9 \\ \left(0.124h^2 - 0.176h + 0.0625\right) = \left(\frac{h}{4.9}\right) \times 4.9 \end{matrix}$$
$$0.6076h^2 - 0.8624h + 0.30625 = h$$

$$0.6076h^2 - 1.8624h + 0.30625 = 0$$

$$h = \frac{1.86 \pm \sqrt{3.46 - 4(0.61)(0.31)}}{2(0.61)}$$

$$h = \frac{1.86 \pm 1.63}{1.22}$$

$$h = 2.9 \text{ m}$$

Expressing without absolute value symbol...

Example:

$$|x+3| \longrightarrow$$

$$|x-5| \longrightarrow \begin{cases} x-5, & \text{if } x-5 \geq 0 \\ -(x-5), & \text{if } x-5 < 0 \end{cases}$$

BBP
 $x \geq 5$
BBN
 $x < 5$

$$|x| = 6$$

$$x = -6 \text{ or } 6$$

$$|-6| = 6$$

$$-(-6) = 6$$

$$|6| = 6$$

$$6 = 6$$

$$|5x+4| \longrightarrow$$

$$5x+4 \geq 0$$

$$5x \geq -4$$

$$x \geq -\frac{4}{5}$$

$$\begin{cases} 5x+4 & \text{if } x \geq -\frac{4}{5} \\ -(5x+4) & \text{if } x < -\frac{4}{5} \end{cases}$$

$$|6|$$

$$|3-4x| \longrightarrow$$

$$3-4x \geq 0$$

$$\frac{-4x \geq -3}{-4 \quad -4}$$

$$x \leq \frac{3}{4}$$

$$\begin{cases} 3-4x & \text{if } x \leq \frac{3}{4} \\ -(3-4x) & \text{if } x > \frac{3}{4} \end{cases}$$

Practice Problems...

Page 363 - 367

#6, 7 c,d , 11, 12, 14, 15, 18, 24

CHAPTER
6

Rational Expressions and Equations

In this chapter, you will learn about the algebra of **rational expressions** and equations. Compare the skills you learn in the chapter with those you learned in the arithmetic of **fractions**. They are very similar.

Definition

A **rational expression** (or **algebraic fraction**) is a fraction with a polynomial in the numerator and a nonzero polynomial in the denominator.

e.g. $\frac{-4}{x}$, $\frac{2y^3 - 4}{-5y + 9}$, $\frac{-4a^3 - 7}{a^2 - 3a}$, $\frac{x - 4}{x^2 + 3x - 28}$

Definition

The **domain of a rational expression (algebraic fraction)** is the set of all real numbers except the value(s) of the variable that result in division by zero when substituted into the expression. To find the values of the variable *excluded* from the domain, **set the denominator equal to zero and solve.**

non-permissible value

- any value for a variable that makes an expression undefined
- in a rational expression, a value that results in a **denominator of zero**
- in $\frac{x+2}{x-3}$, you must exclude the value for which $x-3=0$, which is $x=3$

Examples...

$$\frac{x+2}{3x-5}$$

except..

$$3x-5=0$$

$$3x=5$$

$$x=\frac{5}{3}$$

$$x \in \mathbb{R}, x \neq \frac{5}{3}$$

$$\frac{3x-5}{x^2-13x+30}$$

$$x \in \mathbb{R}, x \neq 3 \text{ or } 10$$

$$\begin{aligned}x^2-13x+30 &= 0 \\(x-10)(x-3) &= 0 \\x &= 10, 3\end{aligned}$$

Example 1

Find the values excluded from domain of each algebraic fraction above.

a. $\frac{-4}{x}$

$x \neq 0$

$$-5y + 9 = 0$$

$$-5y = -9$$

$$y = \frac{9}{5}$$

b. $\frac{2y^3 - 4}{-5y + 9}$

$y \neq \frac{9}{5}$

c. $\frac{-4a^3 - 7}{a^2 - 3a}$

$a \neq 0, 3$

$$a^2 - 3a = 0$$

$$a(a - 3) = 0$$

$$a = 0, 3$$

d. $\frac{x - 4}{x^2 + 3x - 28}$

$x \neq -7, 4$

$$x^2 + 3x - 28 = 0$$

$$(x + 7)(x - 4) = 0$$

$$x = -7, 4$$

Chapter 6

Non-Permissible Values

For each rational expression, determine all non-permissible values. State the domain for each expression. Use the pen tool to write your answers. Drag the Answer tab under each question to reveal the answer.

Answer

1. $\frac{7x}{8x + 16}$

2. $\frac{p^2 - 25}{p^2 - 10p - 24}$

3. $\frac{3}{x^3}$

Hint

4. $\frac{2x - y}{xy}$

$$\begin{aligned}x &\neq 0 \\y &\neq 0\end{aligned}$$

5. $\frac{2x^2 + 5y}{x - y}$

$$x \neq y$$

6. $\frac{2x^2}{x^2 - 4}$

$$\begin{aligned}x^2 - 4 &= 0 \\(x - 2)(x + 2) &= 0 \\x &\neq 2, -2\end{aligned}$$

Simplifying Rational Expressions

Definition

To **simplify a rational expression (an algebraic fraction)** means to write the fraction so that there are no common factors other than 1 or -1.

Steps to simplify an algebraic fraction.

1. Factor the numerator and denominator.
2. Divide out all common factors.

$$\frac{6}{10} = \frac{3}{5}$$

In other words we are reducing algebraic fractions...

- Reduce the following fraction...

$$\frac{8}{10} = \frac{4 \times 2}{5 \times 2} = \frac{4}{5}$$

- Similar process with rational expressions...

$$\frac{5x^2 - 20x}{10x^2 + 20x} = \frac{\cancel{5}(x-4)}{\cancel{10}x(x+2)}$$

$$x \neq 0, -2$$

$$= \frac{1}{2} \frac{(x-4)}{(x+2)}$$

$$= \frac{x-4}{2(x+2)}$$

$$\frac{\cancel{x^2 - x + 7}}{\cancel{x^2 + 2x - 3}}$$

Never cancel
Terms

(Note Monomials are already in factored form)

1. Simplify $\frac{4x^3y^5}{6x^4y^2}$

$$= \frac{2\cancel{y^3}}{3x}$$

2. Simplify $\frac{6x^5y}{12x^2y^3}$

$$= \frac{x^3}{2y^2}$$

3. Simplify $\frac{x^2 - 4}{x^2 - 2x - 8}$

Factor!!
First...

$$\frac{(x-2)(\cancel{x+2})}{(\cancel{x+2})(x-4)}$$

$$= \frac{x-2}{x-4}$$

- diff. of squares
- decomposition
- simple trinomials
- common factor

4. Simplify $\frac{x-4}{4-x} = -1$

$$\frac{10-6}{6-10} = \frac{4}{-4}$$