Homework Questions

12. Amy made the following conjecture: When any number is multiplied

Amy made the following conjecture: When any number is multiplied by itself, the product will be greater than this starting number. For example, in $2 \cdot 2 = 4$, the product 4 is greater than the starting number 2. Meagan disagreed with Amy's conjecture, however, because $\frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4}$ and $\frac{1}{4}$ is less than $\frac{1}{2}$. How could Amy's conjecture be improved? Explain the change(s) you would make.

P.22

1.4

Proving Conjectures: Deductive Reasoning

GOAL

Prove mathematical statements using a logical argument.

Every day, you use deductive thinking to deduce new information.

In this course, you will use this method to deduce the properties of geometric figures and many geometric relationships. For example:

Step A General Statement	And	Step B Particular Statement	Thus	Step C Conclusion
During a game, five players are used on a basketball team.	And	UNB is playing basketball.	Thus	UNB uses five players on the basketball team.
All isosceles triangles have two equal sides.	And	Triangle ABC is isosceles.	Thus	Two sides of Triangle ABC are equal.

In Step A, based on your earlier knowledge (your experience, what you have learned in life) you accept certain general statements to be true. In Step B you are confronted with a particular case that is related to a general statement. Lastly, in Step C, you deduce a conclusion based upon Step A and B.

2

KEY TERMS...

proof

A mathematical argument showing that a statement is valid in all cases, or that no counterexample exists.

deductive reasoning

Drawing a specific conclusion through logical reasoning by starting with general assumptions that are known to be valid.

generalization

A principle, statement, or idea that has general application.

two-column proof

A presentation of a logical argument involving deductive reasoning in which the statements of the argument are written in one column and the justifications for the statements are written in the other column.

We can often use the **transitive property** in deductive reasoning. According to this property, **if two things are equal to the same thing, then they are equal to one another**. We can express this property mathematically:

If a = b and b = c, then a = c.

Mathematical Proofs

A proof is a convincing argument that something is true.

In mathematics, a proof starts with things that are agreed upon, called **postulates** or **axioms**, and then uses logic to reach a conclusion. Conclusions are often reached in geometry by observing data and looking for patterns. As you learned earlier, this type of reasoning is called **inductive reasoning** and the conclusion reached by inductive reasoning is called a **conjecture**.


A proof in geometry consists of a sequence of statements—each supported by a reason—that starts with a given set of premises and leads to a valid conclusion. This type of reasoning is called **deductive reasoning**. Each statement in a proof follows from one or more of the previous statements.

A **reason** (a fact) for a statement can come from the set of given premises or from one of the four types of other premises:

- · Definitions
- Postulates
- · Properties of algebra, equality or congruence
- · Previously proven theorems

Once a conjecture is proved, it is called a **theorem**. As a theorem, it becomes a premise for geometric arguments you can use to prove other conjectures.

Untitled.notebook November 01, 2017



LEARN ABOUT the Math

Jon discovered a pattern when adding integers:

$$\begin{array}{c}
 1 + 2 + 3 + 4 + 5 = 15 \\
 (-15) + (-14) + (-13) + (-12) + (-11) = -65 \\
 (-3) + (-2) + (-1) + 0 + 1 = -5
 \end{array}$$

He claims that whenever you add five consecutive integers, the sum is always 5 times the median of the numbers.



EXAMPLE 1 Connecting conjectures with reasoning

Prove that Jon's conjecture is true for all integers.

Pat's Solution

$$5(3) = 15$$

 $5(-13) = -65$
 $5(-1) = -5$

The median is the middle number in a set of integers when the integers are arranged in consecutive order. I observed that Jon's conjecture was true in each of his examples.

$$210 + 211 + 212 + 213 + 214 = 1060$$

 $5(212) = 1060$

I tried a sample with greater integers, and the conjecture still worked.

Let x represent any integer.

Let S represent the sum of five consecutive integers.

$$S = (x - 2) + (x - 1) + x + (x + 1) + (x + 2)$$

I decided to start my **proof** by representing the sum of five consecutive integers. I chose x as the median and then wrote a generalization for the sum.

proof

A mathematical argument showing that a statement is valid in all cases, or that no counterexample exists.

generalization

A principle, statement, or idea that has general application.

$$S = (x + x + x + x + x) + (-2 + (-1) + 0 + 1 + 2)$$

$$S = 5x + 0$$

I simplified by gathering like terms.

Jon's conjecture is true for all integers.

Since x represents the median of five consecutive integers, 5x will always represent the sum.

Untitled.notebook November 01, 2017

p. 29


EXAMPLE 3 Using deductive reasoning to make a valid conclusion

All dogs are mammals. All mammals are vertebrates. Shaggy is a dog.

What can be deduced about Shaggy?



Oscar's Solution



Transitive Property...

a = b AND b = c therefore, a = c