

OCTOBER 28, 2019

**UNIT 3: SQUARE ROOTS AND
SURFACE AREA**

**SECTION 1.2:
SQUARE ROOTS OF
NON-PERFECT SQUARES**

**K. SEARS
MATH 9**



Oct 28-8:31 AM

WHAT'S THE POINT OF TODAY'S LESSON?

We will continue working on the Math 9 Specific Curriculum Outcome (SCO) "Numbers 4" OR N4 and begin working on "Numbers 6" OR "N6" which state:

N4: "Explain and apply the order of operations, including exponents, with and without technology."

N6: "Determine an approximate square root of positive rational numbers that are non-perfect squares."

Oct 28-8:31 AM



What does THAT mean???


For this unit, SCO N4 means that we will learn how to find the square root (the number that was multiplied by itself) of numbers both with and without a calculator.

SCO N6 means that we will use calculators and "benchmarking" to estimate the square root (the number that was multiplied by itself) of non-perfect squares like 15, 7.5 and $\frac{19}{6}$.



Oct 28-8:31 AM

WARM-UP:



A student has a **rectangular** piece of paper **7.2 cm** by **1.8 cm**. She cuts the paper into parts that can be rearranged to form a **square**.

- a) What is the side length of the **square**?
- b) What are the fewest cuts the student could have made? Explain.



Oct 29-10:21 AM

HOMEWORK QUESTIONS?

(pgs 11 / 12 / 13, #5, 8, 9, 10, 12, 13, 14 and 17)

$$\begin{array}{l|l|l} \sqrt{3.7} & \sqrt{3.6} & \sqrt{0.36} \\ \hline \approx 1.923\dots & \approx 1.89\dots & = 0.6 \\ \text{Not} & \text{Not} & \text{Yes} \end{array}$$

Oct 28-8:56 AM

THE FIRST 20 PERFECT SQUARES:

$1^2 = 1 \times 1 = 1$	$11^2 = 11 \times 11 = 121$
$2^2 = 2 \times 2 = 4$	$12^2 = 12 \times 12 = 144$
$3^2 = 3 \times 3 = 9$	$13^2 = 13 \times 13 = 169$
$4^2 = 4 \times 4 = 16$	$14^2 = 14 \times 14 = 196$
$5^2 = 5 \times 5 = 25$	$15^2 = 15 \times 15 = 225$
$6^2 = 6 \times 6 = 36$	$16^2 = 16 \times 16 = 256$
$7^2 = 7 \times 7 = 49$	$17^2 = 17 \times 17 = 289$
$8^2 = 8 \times 8 = 64$	$18^2 = 18 \times 18 = 324$
$9^2 = 9 \times 9 = 81$	$19^2 = 19 \times 19 = 361$
$10^2 = 10 \times 10 = 100$	$20^2 = 20 \times 20 = 400$

Which numbers are between the perfect squares 1 and 4?
9 and 16? What do you think their square roots will be?

Sep 12-6:48 PM


WARM UP:

Please turn to **page 14** in *MMS9*.

What is the safe height up the wall for the ladder?

Sep 21-9:21 PM


**How could you check if the ladder is safe?
Try to do this without a calculator.**



$$c^2 = a^2 + b^2$$
$$a^2 = c^2 - b^2$$

$$a^2 + b^2 = c^2$$
$$2^2 + b^2 = 9^2$$
$$4 + b^2 = 81$$
$$b^2 = 81 - 4$$
$$b^2 = 77$$
$$\sqrt{b^2} = \sqrt{77}$$
$$b \doteq 8.8 \text{ m}$$



Sep 21-9:39 PM



Pythagoras of Samos
(about 569 BC - about 475 BC)

Pythagoras was a Greek philosopher who made important developments in mathematics, astronomy and the theory of music. The theorem now known as Pythagoras' theorem was known to the Babylonians 1000 years earlier, but he may have been the first to prove it.


Pythagoras discovered a relationship between the areas of the squares drawn on the sides of a right-angled triangle.




Sep 21-11:28 PM

PYTHAGOREAN THEOREM:

We know that a right triangle is a triangle containing a 90° angle.




Hypotenuse
(always the longest side)



Sep 21-11:29 PM


Pythagoras found out that when you have a right triangle, $leg^2 + leg^2 = hypotenuse^2$

$$a^2 + b^2 = c^2$$



Sep 21-11:31 PM

Estimate how far up a wall a 6.1 m long ladder will reach if its base is 1.5 m from the wall.



$$\begin{aligned}
 a^2 &= c^2 - b^2 \\
 &= 6.1^2 - 1.5^2 \\
 &= 37.21 - 2.25 \\
 &= 34.96 \\
 a &= \sqrt{34.96} \\
 &\doteq 5.91
 \end{aligned}$$

$$\begin{aligned}
 a^2 + b^2 &= c^2 \\
 1.5^2 + b^2 &= 6.1^2 \\
 2.25 + b^2 &= 37.21 \\
 b^2 &= 37.21 - 2.25 \\
 b^2 &= 34.96 \\
 \sqrt{b^2} &= \sqrt{34.96} \\
 b &\doteq 5.9 \text{ m}
 \end{aligned}$$


Sep 21-9:41 PM

We will learn two strategies for estimating the square root of a decimal number that is not a perfect square:

- 1) benchmarks
- 2) using a calculator

Sep 21-10:00 PM

BENCHMARKS: Estimate $\sqrt{7.5}$.



Sep 21-10:03 PM

USING A CALCULATOR: Estimate $\sqrt{7.5}$.

$$\sqrt{7.5} \doteq 2.738\ 612\ 788$$

$$\begin{aligned}\text{Check: } & 2.738\ 612\ 788^2 \\ & = 7.500\ 000\ 003\end{aligned}$$

Since this number is not EXACTLY equal to 7.5, the square root is an approximation.

Sep 21-10:05 PM

Let's look at the 4 examples on pages 15 and 16 of *MMS9* for estimating a square root of a fraction using both the benchmark and calculator methods:

Sep 21-10:14 PM

a) $\sqrt{8/5}$


$= \sqrt{\frac{9}{4}}$

$= \frac{3}{2}$

$= 1.5$


1.26 actual

b) $\sqrt{3/10}$



Sep 21-10:21 PM

c) $\sqrt{3/7}$



$= \sqrt{\frac{3}{7}}$

$= \sqrt{0.5}$


$= \sqrt{0.49}$

$= 0.7$

d) $\sqrt{19/6}$

$= 3.\overline{16}$

Sep 21-10:24 PM



Oct 28-1:36 PM

Let's look at the example on page 17 of *MMS9* for finding a number with a square root between two given numbers:

Sep 21-10:27 PM

Identify a decimal number that has a square root between 10 and 11. Check the answer.

Method 1:

$$10^2 = 100$$

$$11^2 = 121$$


Any number between 100
and 121 has a square root
between 10 and 11.

Method 2:

One decimal number
between 10 and 11 is
10.4 .

Sep 21-10:34 PM

Let's practice some more with the Pythagorean Theorem. Calculate the area of carpet needed to cover this ramp.



$$A = l \times w$$

$$\begin{aligned} c^2 &= a^2 + b^2 \\ &= 6.5^2 + 1.5^2 \\ &= 42.25 + 2.25 \\ &= 44.50 \end{aligned}$$

$$\begin{aligned} c &= \sqrt{44.5} \\ &= 6.67 \text{ m} \end{aligned}$$

$$\begin{aligned} A &= l \times w \\ &= (2.2)(6.67) \\ &= 14.674 \text{ m}^2 \end{aligned}$$

Sep 22-8:07 AM

***MMS9*, page 18: DISCUSS THE IDEAS**

1.

2.

3.

Sep 21-10:44 PM

CONCEPT REINFORCEMENT:

MMS9

Page 18: #4, 5 (change to decimal numbers first), 6 and 7

Page 19: #9 (use your calculator), 10, 11, 13, 14 and 16

Page 20: #17, 19, 20 and 21

Sep 12-9:56 PM

MID-UNIT REVIEW:

MMS9

Page 21: #1 to #11

Sep 21-11:26 PM